


10/516600

PCT/EP 03/06194



INVESTOR IN PEOPLE



The Patent Office  
Concept House  
Cardiff Road  
Newport  
South Wales  
NP10 8QQ

03 DEC 2004

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.


BEST AVAILABLE COPY

Signed

Dated

*R. Mahoney*  
27 March 2003

**PRIORITY DOCUMENT**  
SUBMITTED OR TRANSMITTED IN  
COMPLIANCE WITH  
RULE 17.1(a) OR (b)



The  
**Patent  
Office**

1/77  
14JUN02 E725681-1 000524  
P0177700 0.00-0213612.5

**Request for grant of a patent**

(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)

The Patent Office

Cardiff Road  
Newport  
Gwent NP10 8QQ

1.	Your reference	N-32529P1		
2.	Patent application number (The Patent Office will fill in this part)	0213612.5		13 JUN 2002
3.	Full name, address and postcode of the or of each applicant (underline all surnames)	NOVARTIS NUTRITION AG MONBIJOUSTRASSE 118 3001 BERN SWITZERLAND  Patent ADP number (if you know it) If the applicant is a corporate body, give the country/state of its incorporation		
4.	Title of invention	Organic compounds		
5.	Name of your agent (if you have one)  "Address for service" in the United Kingdom to which all correspondence should be sent (including the postcode)	B.A. YORKE & CO. CHARTERED PATENT AGENTS COOMB HOUSE, 7 ST. JOHN'S ROAD ISLEWORTH MIDDLESEX TW7 6NH  Patents ADP number (if you know it)		
6.	If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or of each of these earlier applications and (if you know it) the or each application number	Country	Priority application number (if you know it)	Date of filing (day/month/year)
7.	If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application	Number of earlier application	Date of filing (day/month/year)	
8.	Is a statement of inventorship and of right to grant of a patent required in support of this request? (Answer 'Yes' if: a) any applicant named in part 3 is not an inventor, or b) there is an inventor who is not named as an applicant, or c) any named applicant is a corporate body. (see note (d))	Yes		

7143563001

**Patents Form 1/77**

9. Enter the number of sheets for any of the following items you are filing with this form. Do not count copies of the same document

Continuation sheets of this form

Description 25

Claim(s) 2

Abstract

Drawing(s)

10. If you are also filing any of the following, state how many against each item.

Priority documents

Translations of priority documents

Statement of inventorship and right to grant of a patent (*Patents Form 7/77*)

Request for preliminary examination and search (*Patents Form 9/77*) ONE

Request for substantive examination (*Patents Form 10/77*)

Any other documents (please specify)

11.

I/We request the grant of a patent on the basis of this application

Signature

Date

B. A. Yorke & Co.

B.A. Yorke & Co.

13 June 2002

12. Name and daytime telephone number of person to contact in the United Kingdom

Mrs. E. Cheetham  
020 8560 5847

**Warning**

*After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.*

**Notes**

- If you need help to fill in this form or you have any questions, please contact the Patent Office on 0645 500505.
- Write your answers in capital letters using black ink or you may type them.
- If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.
- Once you have filled in the form you must remember to sign and date it.
- For details of the fee and ways to pay please contact the Patent Office.

### Organic Compounds

The present invention concerns a viscous soluble fiber composition.

The invention also relates to a method for modulating the viscosity of a viscous soluble fiber composition and to a method for preventing or treating diabetes, hypercholesterolemia, obesity, and/or metabolic syndrome in a mammal.

### Background of the Invention

The importance of dietary fiber in the daily diet has been emphasized over the last several years through the recognition of its effects in the prevention and/or treatment of such diseases as hyperlipidemia, coronary heart disease, arteriosclerosis, colon diverticulum, colon cancer, heart diseases, hypercholesterolemia, obesity, hypertension, diabetes or metabolic syndrome.

Metabolic syndrome can be characterized by at least one of the following abnormalities: abdominal obesity, hyperlipidemia, hypercholesterolemia, high blood pressure and high blood glucose level. Recent studies have shown that more than 20 % of the US adult population has the metabolic syndrome.

Soluble fibers act in promoting bowel action. The bulk-forming property of such fiber also contributes to fill the stomach, which has a satiety effect. Soluble fibers delay the emptying of the stomach and slow the transit of chyme in the upper gastrointestinal tract, resulting in slower absorption rates and altered hormonal responses to the absorbed nutrients. It has been proposed that viscous soluble fibers, such as konjac gum or oat beta glucan, may reduce the need for insulin, improve glycemia, reduce serum (LDL) cholesterol, slow the appearance of glucose in the blood and inhibit bile acid absorption in the ileum.

A potential difficulty in adopting viscous soluble fiber in a composition is the viscosity that is generally associated with levels considered to be therapeutic in the treatment of certain diseases, such as diabetes, or metabolic syndrome. In particular, the high viscosity may render problematic the preparation of a composition for oral and enteral administration, especially for patients requiring tube-

feeding. Therefore there is a need for a viscous soluble fiber composition, which is non-viscous, i.e. sufficiently low viscous to be enterally and/or orally administered.

Present inventors have surprisingly found that proteins, such as wheat protein, egg protein, collagen protein, whey protein, casein protein, gluten, pea protein and/or hydrolysates thereof, have the ability to lower the viscosity of viscous soluble fiber compositions, such as konjac gum and/or xanthan gum and/or guar gum and/or beta glucan and/or pectin compositions. According to the invention nutritional compositions comprising high level of high viscous soluble fiber may be formulated, which are convenient to administer, yet without necessarily increasing the glucidic value of such compositions. After ingestion, the viscosity of the soluble fibers comprised in the compositions according to the invention may re-increase because of the digestion and the consecutive hydrolysis of the viscosity-lowering protein.

#### **Summary of the Invention**

In a first aspect of the invention there is provided a composition comprising one or more viscous soluble fiber(s) and one or more viscosity-lowering protein(s).

In another aspect of the invention there is provided a composition according to the invention for use as a medicament.

In a further aspect of the invention there is provided a composition according to the invention for use in meeting the dietary fiber intake of a mammal.

In yet another aspect of the invention there is provided a use of the compositions according to the invention in the manufacture of a medicament or nutritional composition for the prevention or treatment of metabolic syndrome and/or diabetes, and/or for lowering blood cholesterol level.

In another aspect of the invention there is provided a method of treating and/or preventing diabetes and/or metabolic syndrome and/or of lowering blood cholesterol level in a mammal in need of such a treatment comprising administering to said mammal an effective amount of a composition according to the invention.

In a further aspect of the invention there is provided a method for providing viscous soluble fibers to a mammal in need of such fibers comprising administering to said mammal a composition according to the invention.

In another aspect of the invention there is provided a use of protein to modulate the viscosity of a composition comprising one or more viscous soluble fiber(s), wherein said protein is selected from the group consisting of wheat protein, egg protein, collagen, whey protein, casein, soy protein, pea protein, muscle protein, gluten, fibrillar protein, silk protein, hydrolysates thereof and mixtures thereof.

#### **Detailed Description of the Invention**

The term "soluble fiber" as used herein refers to non-starch polysaccharides characterized as being soluble by using the official method of the Association of Official Analytical Chemists (Prosky et al, 1988; J. Assoc. Off. Anal. Chem, 70, 5, 1017), e.g. water soluble fibers, e.g. water soluble at room temperature.

As used herein, the term "viscous fiber" refers to fibers which are able to increase the viscosity of the contents of the stomach and the small intestine and to slow gastric emptying rates. Typically, compositions comprising 1% by weight of such viscous fibers have a viscosity higher than 500 mPa.s (500 centipoises) at room temperature, e.g. temperature from 20°C to 25°C, preferably more than 1000 mPa.s, even more preferably more than 1500 mPa.s. In another aspect of the invention, such viscous fibers have a molecular weight between 20'000 and 5 millions Da, preferably 100'000 and 2 millions Da.

For the purpose of the present invention, the term "compositions of the invention" encompasses compositions comprising viscous soluble fiber and viscosity-lowering protein. Such compositions may be non-viscous, e.g. less viscous than same compositions in which the viscosity-lowering protein is absent, and/or hydrolyzed, e.g. moderately or extensively hydrolyzed. In another aspect of the invention, the compositions according to the invention, e.g. the non viscous compositions of the invention, may be defined by a viscosity less than about 500 mPa.s, preferably less than about 250 mPa.s, more preferably less than about 150 mPa.s, even more preferably less than about 100 mPa.s

and most preferably less than about 60 mPa.s. It has to be understood that such viscosities are measured with a Brookfield LVDV II viscosimeter at room temperature, e.g. temperature from 20° to 25°C, and for a spindle and a shear rate in agreement with the product thickness. For example, viscosities lower than 60 mPa.s are measured with a LVDV II viscosimeter having a spindle of 1 and at 100 rotations per minute (rpm).

The compositions of the invention have the advantageous characteristic that the viscosity may be modulated, for example reversed, delayed, or re-increased. As used herein, "modulated viscosity" refers to the viscosity of a soluble viscous fiber composition that has been modulated, i.e. lowered, or totally or partially re-increased, e.g. by moderately or extensively hydrolyzing the viscosity-lowering protein. As used herein, "delayed viscosity" refers to the viscosity of compositions of the invention which has been re-increased, e.g. by moderately or extensively hydrolyzing the viscosity-lowering protein. The hydrolysis of the viscosity-lowering protein may occur in vivo, i.e. after ingestion of the compositions of the invention, e.g. non-viscous compositions, converting the compositions of the invention from low or non viscous to high viscous compositions, resulting in e.g. a slower glucose absorption and/or in a lower or slower glucose and insulin response to carbohydrate loads. In one aspect of the invention, there is provided a composition comprising viscous soluble fiber(s) and viscosity-lowering protein(s) wherein the viscosity-lowering effect of the protein, e.g. the viscosity of the composition, is reversible.

In one embodiment of the invention, compositions of the invention may consist exclusively or essentially of viscous soluble fiber and viscosity-lowering protein. Alternatively, the compositions of the invention may comprise other nutritional or pharmaceutical components in addition to viscous soluble fiber and viscosity-lowering protein.

Examples of suitable viscous soluble fibers according to the present invention include agar, alginates, carubin, pectin, beta-glucan, such as oat beta-glucan, carrageenans, furcellaran, inulin, arabinogalactan, pectin and its derivatives, cellulose and its derivatives, scleroglucan, psyllium, such as psyllium seed husk, mucilages and gums. In the context of the present invention the term "pectin" refers preferably to pectins from fruits and vegetables, and more preferably to pectins from citrus fruits and apple. The term "carrageenans" as used herein encompasses in particular kappa, lambda and iota carrageenans. According to the invention, gums and mucilages are preferably plant exudates. In

particularly to konjac gum, xanthan gum, guar gum (Guaran gum), locust bean gum, tara bean gum, gum tragacanth, arabic gum, karaya gum, gum ghatti, gellan gum and other related sterculia gum, Alfalfa, Clover, Fenugreek, Tamarind flour. Native and modified, e.g. hydrolyzed, soluble viscous fibers, especially gum arabic, are acceptable for the invention. According to the invention, the preferred viscous fibers are xanthan gum, pectin, guar gum, beta glucan, konjac gum, and any mixtures thereof.

In one embodiment of the invention, the compositions of the invention do not comprise hydrolyzed soluble viscous fiber, e.g. do not comprise extensively hydrolyzed soluble viscous fiber. In another aspect of the invention, the compositions of the invention may comprise native guar gum, xanthan gum, beta glucan and/or konjac gum. Optionally the compositions according to the invention may comprise moderately hydrolyzed soluble viscous fiber, preferably moderately hydrolyzed xanthan gum, beta glucan, konjac gum, guar gum or any combination thereof. As used herein, "moderately hydrolyzed" viscous fiber refers to fibers having a molecular weight between 20'000 and 5 millions Da, preferably between 100'000 and 2 millions Da.

The fiber of the compositions of the invention may consist essentially or exclusively of viscous soluble fibers, optionally in the form of native viscous soluble fibers, or in the form of moderately hydrolyzed viscous soluble fiber. In a preferred embodiment of the invention, the compositions of the invention do not comprise non-viscous soluble fiber. In yet another embodiment of the invention, the compositions of the invention do not comprise inulin.

The compositions of the invention may comprise one viscous soluble fiber, preferably konjac gum, xanthan gum, guar gum, beta-glucan or pectin, even more preferably guar gum. In another embodiment of the invention, the compositions according to the invention may include a mixture of viscous soluble fibers, preferably a mixture of guar gum and one or more viscous soluble fibers selected from the group consisting of pectin, beta-glucan, xanthan gum, arabic gum, carageenan and konjac gum, more preferably a mixture of guar gum and one or more viscous soluble fibers selected from the group consisting of xanthan gum and konjac gum, and even more preferably a mixture of guar gum and xanthan gum.

Xanthan gum, the extracellular polysaccharide from *Xanthomonas campestris* and some related microorganisms, is produced on a nutritive medium comprising glucose, a nitrogen sources and



minerals. It is used in the food industry as a stabilizer or thickener. It is also known to interact synergistically with other hydrocolloids like guar gum. Xanthan gum is commercially available from Satiexane CX2QD, Degussa Texturant System, France.

Konjac glucomannan, also called konjac mannan or konjac gum, has the highest viscosity of any dietary fiber known. Konjac gum has a high capacity for water absorption and can swell to about 200 times its original volume. In food industry, konjac gum is often added to starch because their interaction leads to a considerable increase in the viscosity and to the formation of gel, which can be maintained during cooking. Konjac gum is also known to interact synergistically with hydrocolloids such as kappa-carrageenan, xanthan gum or locust bean gum to form heat-stable gels. Konjac gum may also gel by itself if heated after alkali addition. According to the present invention, konjac gum may be extracted from the *Amorphophallus* species, more particularly from *A. riviera* (often referred to as *A. konjac*), *A. oncophyllus*, *A. variabilis*, *A. bulbifera* and *A. blumeli*, more preferably from *A. riviera*, and even more preferably from *A. riviera* C. Koch. In particular konjac gum is produced from the root of the plant *Amorphophallus*, and more preferably from the tuber thereof. Konjac flour may be prepared by drying, pulverizing and winnowing the plant. Several techniques are known in the art for separating konjac glucomannan from konjac flour. As used herein the term "konjac gum" encompasses konjac glucomannan and konjac flour. The konjac gum is commercially available from Propol RS, Shimizu, Japan.

Guar Gum is obtained from the seed endosperms of the leguminous plant *Cyamopsis tetragonolobus* (Linné) Taub. and consists mainly of a high molecular weight hydrocolloidal polysaccharide, composed of galactose and mannose units combined through glycoside linkages. Guar Gum is used in the food industry as a stabilizer or thickener. It is also known to interact synergistically with other hydrocolloids like xanthan gum. Guar gum is commercially available from Polygum 75, Polygal AG, Switzerland (high molecular weight type) and from Lygomme 3, Degussa Texturant Systems, France (low molecular weight type).

Mixed-linkage (1→3)(1→4)-β-D-glucan (beta-glucan) is the main endospermic cell wall polysaccharide of oats and barley, but also present, in lesser amounts, in rye and wheat. The subaleurone and subembryo layers of dehulled oat are particularly rich in beta-glucan, whereas in the inner starch endosperm concentrations are low. Beta-glucan can therefore be enriched in bran. The present invention provides a process for the preparation of a dietary fiber composition from the bran of oats and barley, which is characterized in that it contains a high concentration of beta-glucan.

Pectin is a purified carbohydrate product obtained from the dilute acid extract of the inner portion of the rind of citrus fruits or from apple pomace. It consists mainly of partially methoxylated polygalacturonic acids. It is commercially available from Riedel de Haen AG, Germany.

Details of the active ingredients of the compositions of the invention may be obtained from the relevant manufacturers, the contents of which are hereby incorporated by reference.

The relative proportion of the active ingredients of the compositions of the invention will, of course, vary considerably depending on the particular type of composition concerned, e.g. whether it is a liquid or solid form, or whether it is provided in nutritional form. All indicated proportions and relative weight ranges described below are accordingly to be understood as being indicative of preferred or individually inventive teaching only and not limiting the invention in its broadest aspect.

According to the present invention, the term "soluble viscous fiber" is referred to as fiber concentrate or not pure fiber source. Fiber concentrates may be pectin and gum, more preferably gum arabic, locust bean gum, guar gum, psyllium seed husk. Fiber concentrate of the invention may comprise more than 90 % by weight of fiber, based on the total dry weight of the fiber concentrate, preferably more than 95% by weight. Not pure fiber source may comprise more than 80% by weight of fiber.

Furthermore the compositions of the invention may comprise insoluble fibers, such as cellulose, hull fibers from legumes and grains, such as pea hull fiber, oat hull fiber, barley hull fiber, soy hull fiber. In one embodiment of the invention, the ratio of soluble viscous fiber to insoluble fiber may be from 0.1 to 10.

The term "viscosity-lowering protein" as used herein refers to proteins or peptides which when added to a viscous soluble fiber composition may lower the viscosity of that composition. According to the present invention these proteins may have a molecular weight lower than about 35 kDa, more preferably lower than about 30 kDa, even more preferably from about 0.2 to about 30 kDa, and even more preferably from about 1 to about 25 kDa. These values refer to an approximate molecular weight range with values falling above and below the given range. The viscosity-lowering proteins of the invention may be in the form of native protein or in form of hydrolysate.

Examples of "viscosity-lowering protein" according to the present invention include wheat protein, soy protein, pea protein, egg protein, collagen, whey protein, casein, gluten, muscle proteins from fish and animals, fibrillar proteins, silk proteins, e.g. keratin, sericin or fibroin, and mixtures thereof. Accordingly, in one aspect the present invention provides compositions comprising wheat protein, egg protein, collagen, whey protein, casein, soy protein, pea protein, fibrillar proteins, silk proteins, e.g. keratin, sericin or fibroin, hydrolysates thereof or mixtures thereof, preferably wheat protein, egg protein, collagen, pea protein, fibrillar proteins, silk proteins, e.g. keratin, sericin or fibroin, hydrolysates thereof or mixtures thereof. In another aspect the present invention provides compositions of the invention comprising whey protein, casein and/or hydrolysate thereof, with the proviso that said compositions do not comprise inulin.

In another aspect of the invention, there is provided a composition of the invention comprising soy protein and/or hydrolysate thereof, with the proviso that said composition does not comprise phytoestrogen.

In yet another aspect of the invention, there is provided a composition of the invention comprising viscous soluble fiber selected from the group consisting of xanthan gum, arabic gum, guar gum, pectin, beta-glucan, and mixtures thereof, with the proviso that said compositions do not comprise inulin.

In a further embodiment of the invention, the compositions of the invention may comprise a viscous soluble fiber selected from the group consisting of gum, mucilage, carrageenan, and mixtures thereof, preferably a viscous soluble fiber selected from the group consisting of konjac gum, locust bean gum, tara bean gum, gum tragacanth, karaya gum and mixtures thereof.

In a preferred embodiment of the invention, the compositions of the invention may comprise guar gum, xanthan gum, konjac gum, pectin and/or beta-glucan in combination with a viscosity-lowering protein selected from the group consisting of wheat protein, egg protein, collagen, pea protein, fibrillar proteins, silk proteins, e.g. keratin, sericin or fibroin, hydrolysates thereof and mixtures thereof.

In another preferred embodiment of the invention, the compositions of the invention consist essentially of or exclusively of guar gum and collagen hydrolysate.

In another aspect of the invention, there is provided a composition of the invention comprising a viscosity-lowering protein which has been hydrolyzed, e.g. a protein not further able to decrease the viscosity of the composition. As used herein, "hydrolyzed protein" refers to protein which has a molecular weight lower than about 0.5 kDa, more preferably lower than 0.2 kDa, and even more preferably lower than about 0.1 kDa. It will be appreciated that according to the invention one or a mixture of proteins may be used.

The amount of viscous soluble fiber incorporated into the compositions of the invention may depend on the nature and/or the molecular weight of the viscous fiber, the nature and the amount of the viscosity-lowering protein, the form of the compositions of the invention, i.e. a powder or a composition ready-for-consumption, -e.g. ready-to-drink compositions or instant drink-, or on the final viscosity to be reached, i.e. the type of product in which the compositions will be incorporated. Typically, the more viscous the viscous soluble fiber is, the less it amounts in the compositions.

Accordingly, suitable amounts of viscous soluble fiber comprised in ready-for-consumption compositions according to the invention are in the range of up to about 40% by weight, preferably up to about 20% by weight, more preferably up to about 15% by weight, even more preferably from about 0.05 to about 10% by weight, more preferably from about 0.1 to about 5% by weight, most preferably from about 0.5 to about 2.5% by weight, and even more preferably about 1% by weight, based on the total weight of the ready-for-consumption composition.

The compositions of the invention in powder form may comprise viscous soluble fiber in an amount of about 0.5 to about 60% by weight, more preferably in an amount of about 4 to about 40% by weight, more preferably in an amount of about 10 to about 30% by weight and even more preferably in an amount of about 8 to about 20% by weight, based on the total weight of the powder composition.

In one preferred embodiment of the invention, the compositions according to the invention, the weight ratio of xanthan gum to konjac gum is from about 0.5 to about 10, more preferably from about 1 to about 5, and even more preferably from about 2 to about 4. In another embodiment of the invention, the weight ratio of guar gum to xanthan gum of the compositions of the invention is from about 0.01 to about 100, more preferably from about 0.05 to about 10, and even more preferably from about 0.1 to about 5.

The suitable amount of viscosity lowering-proteins incorporated into the compositions of the invention may depend on the molecular weight and/or on the nature of the protein, for example on its three-dimensional structure, e.g. globular, fibrillar, denaturated structure, its charge, its pHi.

The amount of viscosity-lowering protein incorporated into the ready-for-consumption compositions of the invention may vary from about 1 to about 60% by weight, preferably from about 2 to about 20 % by weight, more preferably from about 3 to about 10% by weight, even more preferably from about 4 to about 8 % by weight, based on the total weight of the composition.

In one embodiment of the invention, the compositions of the invention in powder form comprise an amount of viscosity-lowering protein from about 0.05 to 90% by weight, preferably from about 0.5 to about 80% by weight, more preferably from about 20 to about 60% by weight of the composition, based on the total weight of the composition.

In another embodiment of the invention, the weight ratio of viscous soluble fiber to viscosity lowering-protein in the compositions of the invention, in powder or in ready for consumption form, may be from about 0.01 to about 20, preferably from about 0.05 to about 10, more preferably from about 0.1 to about 1, and most preferably from about 0.1 to about 0.4.

In one aspect of the invention, the total amount of viscous soluble fiber and viscosity lowering-protein in the compositions of the invention, in powder or in ready for consumption form, may be from about 1 to about 65% by weight, more preferably from about 1 to about 30% by weight, even more preferably from 2 to 15% by weight and most preferably from about 5 to about 15% by weight, based on the total weight of the composition.

In one embodiment of the invention, the compositions of the invention may comprise other "proteinaceous material" in addition to the viscosity lowering-protein. "Proteinaceous material" as used herein refers to any form of protein, peptide, free amino acid, mixtures of amino acids, and combinations thereof.

In a further embodiment of the invention, the compositions of the invention may comprise, in addition to soluble viscous fiber and viscosity lowering-protein, other nutritional components like fat,

carbohydrate, minerals and vitamins, for example, among others, calcium, magnesium, iron, zinc, phosphorus, vitamin D and/or vitamin K.

In another embodiment of the invention, the carbohydrate content of the compositions according to the invention, including viscous soluble fiber, may be less than about 60 en%, preferably less than about 55 en%, more preferably less than about 30 en%, most preferably less than about 20 en%, based on the total caloric content of the composition.

In yet another embodiment of the invention, the compositions according to the invention may not comprise starch-derived carbohydrates, such as maltodextrin, modified starch, amylose starch, topioca starch, corn starch, or mixtures thereof.

In another embodiment of the invention, the compositions according to the invention may not comprise saccharose.

In a further embodiment of the invention, the compositions according to the invention may comprise fructose and/or galactose.

Further components of the compositions according to the invention may include any bioactive compounds or extracts which are known to have health benefits, especially for lowering blood cholesterol level in mammals, such as statins, bile acid sequestrants, nicotinic acid or fibric acids, and/or for treating diabetes, such as sulfonylureas, biguanides, alpha-glucosidase inhibitors, thiazolidinediones, meglitinides or D-phenylalamine. For example, the composition of the invention may be provided in the form of a kit for separate, sequential or simultaneous administration in conjunction with such medicines for lowering blood glucose and/or blood cholesterol level. These medicines may conveniently be formulated together with viscous soluble fiber, such as konjac gum, xanthan gum, beta-glucan or guar gum, and a viscosity-lowering protein in standard pharmaceutical dosage forms.

In one aspect of the present invention, the compositions according to the invention can readily be incorporated into pharmaceutical or nutritional formulations, typically nutraceuticals, dietary supplements, functional food and beverage products.

In a further aspect of the invention, the compositions of the invention may be used as a medicament. Accordingly the compositions of the invention may be administered in pharmaceutical form or as a dietary supplement, preferably in combination with at least one pharmaceutically or nutritionally acceptable carrier.

In a yet further aspect of the invention, there is provided a medicament, nutritional or pharmaceutical formulation, for example dietary supplement, comprising the composition of the invention. The medicament, nutritional or pharmaceutical composition of the invention may optionally comprise pharmaceutical acceptable carriers. Further, according to the invention there is provided a combined pharmaceutical preparation for simultaneous, separate or sequential use for the treatment or prevention of diabetes comprising soluble viscous fiber and viscosity-lowering protein and one or more drug(s) for lowering blood glucose and/or one or more drug(s) for lowering blood cholesterol level.

Pharmaceutical compositions and dietary supplements may be provided in the form of soft gel, sachets, powders, syrups, liquid suspensions, emulsions and solutions in convenient dosage forms. In soft capsules the active ingredients are preferably dissolved or suspended in suitable liquids, such as fatty oils, paraffin oil or liquid polyethylene glycols. Optionally stabilisers may be added.

Oral pharmaceutical or dietary supplement forms may be made by conventional compounding procedures known in the pharmaceutical art, that is, by mixing the active substances together with edible pharmaceutically acceptable solid or liquid carriers and/or excipients, e.g. fillers such as cellulose, lactose, sucrose, mannitol, sorbitol, and calcium phosphates and binders, such as starch, gelatin, tragacanth, methylcellulose and/or polyvinylpyrrolidone (PVP). Optional additives include lubricants and flow conditioners, e.g. silicic acid, silicon dioxide, talc, stearic acid, magnesium/calcium stearates, polyethylene glycol (PEG) diluents, disintegrating agents, e.g. starch, carboxymethyl starch, cross-linked PVP, agar, alginic acid and alginates, colouring agents, flavouring agents, and melting agents. Dyes or pigments may be added to the tablets or dragée coatings, for example for identification purposes or to indicate different doses of active ingredient.

Optionally, the compositions according to the invention may be nutritionally complete, i.e. may include vitamins, minerals, trace elements as well as nitrogen, carbohydrate and fatty acid sources so that they may be used as the sole source of nutrition supplying essentially all the required daily amounts of vitamins, minerals, proteins, lipids, carbohydrates and trace elements.

compositions of the invention may be provided in the form of a nutritionally balanced complete meal, e.g. suited for oral or tube feeding.

Alternatively, the compositions of the invention may be provided as part of a meal, i.e. a nutritional supplement, e.g. in the form of a health drink.

In another aspect of the invention, there is provided a method to meet the dietary fiber intake of a mammal, comprising administering to a mammal in need thereof a composition of the invention.

It may be desirable to provide the soluble viscous fiber in the form of a low calorie meal replacement or other nutritional product. In this case the meal replacement or other nutritional product is preferably low fat, i.e. less than about 10 en%, or substantially fat-free, i.e. less than about 2.5 en% contributed by fat, such as about 2 en% fat, based on the total caloric content of the composition. Suitably, a single serving of a low calorie meal replacement will have a caloric value of less than about 1000kcal, and preferably between about 200kcal and about 500kcal. Suitable low calorie nutritional product may include soft drink, such as juice, smoothie or soy-based drink, or dispersed in foods of any sort, such as, dairy bars, soups, breakfast cereals, muesli, candies, tabs, cookies, biscuits, crackers, such as a rice crackers, and dairy products, such as milk-shake, yoghurt drink.

The compositions of the invention optionally comprise conventional food additives, such as any of emulsifiers, stabilizers, sweeteners, flavourings, colouring agents, preservatives, chelating agents, osmotic agents, buffers or agents for pH adjustment, acidulants, thickeners, texturisers, and so on.

In a further aspect of the invention, there is provided a use of compositions of the invention as food additive.

The method of modulating, i.e. increasing or decreasing, the viscosity of the compositions of the invention may be advantageous in the preparation of food products and beverages. For example, the compositions of the invention, e.g. low viscous compositions, may be used for preparing a high viscous nutritional product. The compositions of the invention may also be used to modify the viscosity, to stabilize and/or to texturize food and beverage products, such as ice creams, candies, chocolates, muesli, cookies, biscuits, crackers, chewing-gums, dairy products, desserts, milk-shakes, yogurt drinks, soups and the like. Optionally the compositions of the invention may be used as a fat



substitute in a variety of reduced fat, low fat and fat free foods and beverages, such as cakes, pudding type desserts, butter, peanut butter, cheeses, salad dressings, sauces, margarine, cream cheese and other spreads, snack dips, mayonnaise, sour cream, yogurt, ice cream, frozen desserts, fudge and other confections, and skim milk. The compositions of the invention are also useful, for example as a shortening, in baked goods such as cakes, pies, brownies, cookies, breads, noodles, snack items, such as crackers, graham crackers and pretzels, and similar products.

Suitable product formats according to the present invention include solution, ready-for-consumption composition, e.g. ready-to-drink compositions, instant drink, liquid comestibles, like soft drinks, juice, sports drinks, milk drinks, milk-shakes, yogurt drinks or soup. In a further embodiment of the invention, the compositions of the present invention may be manufactured and sold in the form of a concentrate, a powder, or granules, e.g. effervescent granules, which are diluted with water or other liquid, such as milk or fruit juice, to yield a ready-for-consumption composition, e.g. ready-to-drink compositions or instant drink.

Enteral administration of the compositions of the invention, and preferably oral administration are covered by the present invention.

The amount and dosage regimen of the compositions of the invention to be administered is determined in the light of various relevant factors including the purpose of administration, the age, sex and body weight of individual subject and the severity of the subject's symptoms.

When the composition according to the invention is supplied in the form of a food or beverage, a suitable serving size of the composition may be from about 20 to about 500g, preferably from about 30 to about 300g. If provided in pharmaceutical form, suitable daily doses of the composition of the invention are up to about 250g, preferably 150g, and optimally in the range of about 10 to about 100g. In terms of body weight, an appropriate daily dosage range is about 2 to about 15g composition of the invention per kg. The daily dosage may correspond to a single unit dosage, or may be provided through multiple unit dosages. The exact amounts of the compositions according to the invention may vary between wide limits and may be readily determined in conventional manner.

The compositions of the invention, e.g. medical or nutritional formulations comprising the compositions of the invention, have the characteristic of decreased viscosity when

allowing delivery of an effective amount of viscous soluble fibers. Suitable daily doses of viscous soluble fiber according to the invention are in the range of about 2 mg to about 50 g, preferably in the range of about 25 mg to about 40 g more preferably in the range of about 50 mg to about 30 g and most preferably in the range of about 2 g to about 20 g.

The compositions of the invention may be administered under the supervision of a medical specialist, or may be self-administered.

The compositions of the invention may be administered daily, weekly or monthly, or may be administered annually or even for longer periods in order for example to constantly control the glucose or cholesterol blood level of a patient suffering from diabetes or high blood cholesterol level. Suitable daily dosage regimen may include single or multiple servings per day. Optimally, the composition of the invention is consumed at least once a day on a regular basis, prior to, i.e. pre or post-prandial administration, or during a meal. Preferably, the compositions of the invention may be consumed prior to a meal. In one embodiment of the invention, the composition of the invention may be taken until for example a blood glucose level of 180 mg/dL or less, 2 hours after meals, has been resumed, or until a blood cholesterol level of 200 mg/dL or less has been obtained, and/or a blood LDL cholesterol level of less than 160mg/dL or less than 130mg/dL has been resumed. Since these formulations are safe to consume, subjects with diabetes, metabolic syndrome and/or hypercholesterolemia can continue taking composition according to the invention for as long as required, and preferably until normal blood glucose level, 2 hours after meals, and/or cholesterol level has been resumed.

Pharmaceutical, food or beverage incorporating compositions according to the invention can be safely-consumed by anyone, and are especially recommended for anyone perceived to be at risk from hypercholesterolemia, diabetes or metabolic syndrome, such as elderly, smokers and/or anyone suffering from overweight, obesity, heart disease or high blood pressure. The compositions of the invention may be particularly suitable for subjects suffering from metabolic syndrome, blood sugar disorders, arteriosclerosis, hypertension, hypercholesterolemia, hyperlipidemia, diabetes, intestinal or digestive troubles, Crohn's disease, intestinal diverticula, carcinoma of colon. By lowering blood glucose level, the compositions of the invention have also the effect of counteracting the sequelae of heart disease, high blood pressure, stroke, blindness, blood vessel disease, kidney failure, amputation and nerve damage. "High blood pressure" as used herein refers to a blood pressure persistently

exceeding 140/90 mmHg (systolic/diastolic). By lowering blood cholesterol level, the compositions of the invention may also prevent heart diseases, e.g. heart attacks.

In one embodiment of the invention, the invention pertains to a method of treating and/or preventing metabolic syndrome and/or diabetes and/or hypercholesterolemia, in a mammal, including human, in need of such a treatment, comprising administering to said mammal an effective amount of a composition according to the invention. As used herein, the term "an effective amount" refers to an amount effective to achieve a desired therapeutic effect, such as treating and/or preventing diabetes and/or metabolic syndrome and/or hypercholesterolemia.

In another aspect of the invention, there is provided a method of lowering blood cholesterol level and/or blood glucose in a mammal, including human, in need of such a treatment comprising administering to said mammal an effective amount of a composition according to the invention. Hereinafter the terms "lowering blood glucose" refer to lowering the blood glucose to a level of 180 mg/dL or less 2 hours after meals. As used herein, "lowering blood cholesterol level" refers to a blood cholesterol lowered to a level of less than 240mg/dL, more preferably less than 200 mg/dL, or to a blood LDL cholesterol lowered to less than 190mg/dL, more preferably less than 160 mg/dL, even more preferably less than 130 mg/dL.

In another embodiment of the invention, the compositions according to the invention may be used in the manufacture of a medicament or nutritional formulation for the prevention or treatment of metabolic syndrome and/or diabetes and/or hypercholesterolemia in mammal, including human.

In a further embodiment of the invention, the compositions according to the invention may be used in the manufacture of a medicament or nutritional formulation for lowering blood glucose and/or blood cholesterol level in mammal.

According to the invention, the compositions of the invention may be used in meeting the dietary fiber intake of a mammal.

The method of treatment according to the invention may be combined with diabetes medicines, such as sulfonylureas, biguanides, alpha-glucosidase inhibitors, thiazolidinediones, meglitinides or D-phenylalanine, and with lipid-lowering medicines, such as statins, bile acid sequestrants,

nicotinic acid and fibric acids. According to the invention, a low viscous composition comprising viscous soluble fiber may be produced by admixing such a viscous soluble fiber and a viscosity-lowering protein selected from the group consisting of wheat protein, egg protein, collagen, whey protein, casein, soy protein, pea protein, muscle protein, fibrillar protein, silk proteins, e.g. keratin, sericin or fibroin, gluten and mixtures thereof, more preferably from the group consisting of wheat protein, egg protein, collagen, whey protein, casein and mixtures thereof, in an aqueous medium under conditions sufficient to form a non-viscous composition, e.g. a composition which has a viscosity lower than the viscosity of the same composition in which the viscosity-lowering protein is absent. Suitable examples of aqueous medium include, but are not limited to, water, milk, water-based beverages, milk-based beverages, carbonated beverage, non-carbonated beverage, beer, wine, soy milk and fruit-based beverage, like orange juice. Such aqueous medium may also include specific buffers, like phosphate buffer.

According to the invention, the ingredients may be added simultaneously or sequentially, i.e. the viscosity-lowering protein followed or preceded by the viscous soluble fiber, to the aqueous medium, as defined above.

In one embodiment of the invention, the ingredients of the compositions of the invention may be added in dry form, i.e. as a dry blend of viscous soluble fiber and viscosity-lowering protein. Such a dry blend may be dispersed into an aqueous medium by stirring or mixing, preferably using cold, hot or boiling aqueous medium, and more preferably using aqueous medium at a temperature of about 5°C to about 35°C.

In another embodiment of the invention, the dried ingredients of the compositions of the invention may be hydrated separately before being mixed, i.e. a dispersion of soluble viscous fiber and/or a dispersion of viscosity-lowering protein may be prepared separately and then mixed. Suitable techniques of hydrating viscous soluble fibers and obtaining such dispersions are known in the art. The conditions chosen, such as temperature or stirring time, for hydrating the viscous soluble fiber will depend in particular on the nature and concentration of the fiber. Such conditions are known in the art. To obtain a hydrated soluble viscous fiber dispersion, the soluble viscous fiber may be added to a cold, hot or boiling aqueous medium, more preferably to an aqueous medium at a temperature from about 70°C to about 80°C, and stirred for several hours, for example for 1 hour. Preferably, the concentration of the soluble viscous fiber in the dispersion will vary from about 0.05 to about 15 % by

weight, more preferably from about 0.5 to about 5% by weight, based on the total weight of the dispersion. In one embodiment of the invention, the soluble viscous fiber dispersion may be prepared in a phosphate buffer, e.g. at a pH from about 6.7 to about 7.2. Lower pH values may also be used, i.e. a pH from about 3.8 to about 5.7. For example, a konjac gum dispersion may be prepared by adding konjac flour in an amount of from about 0.5 to about 5% by weight, based on the total weight of the dispersion, to a phosphate buffer solution, having a pH from about 6.7 to about 7.2, at room temperature, and by stirring for approximately 5 minutes. The hydrated viscosity-lowering protein dispersion may be prepared by hydrating the dry viscosity-lowering protein in an aqueous medium at room temperature. The concentration of the viscosity-lowering protein in the dispersion may vary from about 0.1 to about 60% by weight, preferably from about 0.5 to about 35%, more preferably from about 1 to about 10% by weight, based on the total weight of the dispersion. For example, collagen hydrolysate dispersion may be prepared by stirring for less than 5 minutes in a cold aqueous medium.

In a further embodiment of the invention, the dried soluble viscous fiber may be added to a viscosity-lowering protein dispersion, e.g. hydrated protein, or the dried viscosity-lowering protein may be added to a soluble viscous fiber dispersion. In either case, the dried component is added to a dispersion at a low, room or high temperature, for example from about 5°C to about 60°C, more preferably from about 15°C to about 25°C.

According to the invention, the concentration of soluble viscous fiber may vary from about 0.05 to about 30% by weight, preferably from about 0.1 to about 10% by weight, more preferably from about 0.5 to about 5% by weight, and even more preferably from about 0.75% to about 2% by weight, based on the total weight of the composition. The concentration of the viscosity-lowering protein may be from about 1 to about 60% by weight, preferably from about 2 to about 20% by weight, more preferably from about 3 to about 10% by weight, even more preferably from about 4 to about 8% by weight, based on the total weight of the composition. In a further step, the composition of the invention may be dehydrated to obtain a powder.

It will be appreciated that the suitable method of dispersion will be chosen based on the desired properties and/or uses of the compositions of the invention.

According to the invention, the viscosity of the compositions of the invention may be modulated, for example to increase or decrease the viscosity of the composition. One method of a method is a suitable

and/or delay, preferably re-increase, the viscosity of the non-viscous compositions of the invention, by modulating and/or delaying the degradation state of the viscosity-lowering protein. The more viscosity-lowering protein is degraded, i.e. hydrolyzed, the less is its effect on lowering the viscosity, and the more the viscosity re-increases. Suitable methods of hydrolyzing proteins are known in the art. For example, the viscosity-lowering proteins of the compositions of the invention may be hydrolyzed by proteases, such as pepsin, trypsin, chymotrypsin, exopeptidase or endopeptidase, at a temperature from about 20°C to about 60°C, preferably at about 37°C. Optionally, the viscosity of the compositions of the invention, e.g. non viscous or low viscous composition, may be re-increased by heating and simultaneously using an alkaline treatment, e.g. at a pH of about 12, or more preferably by using an acidic treatment, e.g. at a pH of about 4.5, more preferably at a pH below 2. Various degrees of modulation of the viscosity of the compositions of the invention may be obtained by varying the degree of hydrolysis, i.e. the hydrolysis conditions, such as for example temperature, incubation period, nature of enzyme and/or pH. A combination of two or more hydrolysis methods, e.g. of enzymatic and acid hydrolysis, may also be used. In one embodiment of the invention, the composition of the invention may be incubated in the presence of a protease under conditions and time sufficient to hydrolyze the viscosity-lowering protein to the extent that it no longer suppresses the viscosity of the viscous soluble fiber.

In a further other aspect of the invention, there is provided a use of means for hydrolyzing the viscosity-lowering protein(s) to re-increase or reverse the viscosity of a composition comprising viscous soluble fiber(s) and viscosity-lowering-protein(s). According to the invention, such means may include proteases, such as pepsin, trypsin, chymotrypsin, exopeptidase or endopeptidase, as well as acidic or basic solutions in amount sufficient to decrease or increase, respectively, the pH of the composition of the invention, in order to induce hydrolysis of the viscosity-lowering-protein(s).

The invention is further illustrated with reference to the following examples.

### Examples

The konjac gum is from Propol RS, Shimizu, Japan.

The collagen hydrolysate is from Collagel A, Deutsche Gelatine-Fabriken Stoess & Co GmbH, Germany.

High MW Guar type is from Polygum 75, Polygal AG, Switzerland

Low MW Guar type is from Lygomme 3, Degussa Texturant Systems, France.

Xanthan gum is from Satiaxane CX2QD, Degussa Texturant System, France.

Aspartame is from Searle, USA (Nutrasweet™).

Acesulfam K is from Nutrinova, Germany (Sunett™).

**Example 1: Influence of viscosity-lowering protein on a 1% konjac gum composition.**

The konjac gum and the viscosity-lowering protein are admixed together and dispersed in water at temperature of about 25°C. A magnetic agitation is used as dispersion tools and the mixing is kept for 2min. After 1 hour standing time the viscosity is measured.

Protein	Molecular Weight (kDa)	Protein Content (1)	Konjac Content (1)	Protein Solution Viscosity*	Final Solution Viscosity**
Glycin	0.1	10	1	<10	>5000
Wheat	<1	10	1	<10	200
Egg	3-10	10	1	50	250
Collagen	10-18	10	1	10	60
Whey	10-18	10	1	25	500
-	-	-	1	-	>5000

(1) % by weight, based on the total weight of the composition.

\* Viscosity (mPa.s) measured with a Brookfield viscometer (LVDV number 1 spindle, 60 rpm).

\*\* Viscosity (mPa.s) measured with a Brookfield viscometer (LVDV number 4 spindle, 60 rpm) after 30 minutes of standing time.

**Example 2: Influence of the collagen concentration on the viscosity of a 1% konjac gum composition.**

Compositions comprising konjac gum and collagen hydrolysate at different concentrations were prepared as described in Example 1, and their viscosities were measured at 25°C with a Brookfield viscometer using a LVDV number 4 spindle and a shear rate of 60 rpm.

Konjac gum (1)	1	1	1	1	1
Collagen (1)	10	5	5	5	0

(1) % by weight, based on the total weight of the composition.



**Example 3: Influence of collagen on the viscosity of konjac gum and/or xanthan gum compositions.**

The viscous soluble fiber compositions were prepared as described in Example 1, and their viscosities were measured.

Konjac gum <sup>(1)</sup>	-	0.8	0.8	-	1
Xanthan gum <sup>(1)</sup>	0.2	0.2	0.2	0.2	-
Collagen <sup>(1)</sup>	-	-	5	5	-
Viscosity (mPa.s)	120**	1500*	400**	15**	7000*

<sup>(1)</sup> % by weight, based on the total weight of the composition.

\* Viscosity measured with a Brookfield viscometer using a LVDV number 4 spindle and a shear rate of 60 rpm.

\*\* Viscosity measured with a Brookfield viscometer using a LVDV number 2 spindle and a shear rate of 60 rpm.

**Example 4: viscosity of guar gum compositions.**

The compositions were prepared as described in the Example 1. The viscosities were measured with a Brookfield viscometer with LV spindle at 20°C.

**A. Influence of molecular weight and concentration of guar gum on the viscosity.**

High MW guar gum	1	2	3	4	-	-	-	-	-
(g)									
Low MW guar gum (g)	-	-	-	-	1	2	3	4	5
Water (g)	199	198	197	196	199	198	197	196	195
Viscosity (mPa.s)	330	4100	28000	170000	75	1020	4500	15000	58000

**B. Viscosity lowering effect of collagen on guar gum compositions.**

High MW guar gum	3.0	3.0	3.0	-	-	-
(g)						
Low MW guar gum (g)	-	-	-	4.3	4.3	4.3
Collagen (g)	-	10	15	-	10	15
Water (g)	197	187	182	195.7	185.7	180.7
Viscosity (mPa.s)	10000	110	10	10000	110	35

C. Viscosity lowering effect of different collagen concentration on the viscosity of a 1.5% by weight, guar gum composition (high MW type).

Guar gum (g)	3	3	3	3	3	3
Collagen (g)	0	2.5	5	7.5	10	15
Water (g)	197	194.5	192	189.5	187	182
Viscosity (mPa.s)	31000	26000	5300	110	67	29

**Example 5: Viscosity of xanthan gum compositions.**

The viscous soluble fiber compositions were prepared as described in Example 1, and the viscosities were measured with a Brookfield, spindle LV DV at 20°C.

A. Influence of collagen concentration on the viscosity of a 1% by weight xanthan gum composition.

Xanthan gum (g)	2	2	2	2	2	2
Collagen (g)	0	2.5	5	7.5	10	15
Water (g)	198	195.5	193	190.5	188	183
Viscosity (mPa.s)	760	520	3050	6300	30400	41000

B. Influence of collagen on the viscosity of xanthan gum compositions.

Xanthan gum (g)	0.4	0.4	1	1	2	2
Collagen (g)	-	10	-	10	-	10
Water (g)	199.6	189.6	199	189	198	188
Viscosity (mPa.s)	120	15	300	2500	760	30400

C. Influence of the acidity on the viscosity of a 0.5% by weight xanthan gum composition.

Xanthan gum (g)	1	1	1	1
Collagen (g)	5	10	5	10
Citric acid (g)	-	-	0.05	0.05
Water (g)	194	189	194	189
pH	5.55	5.55	5.15	5.15

Viscosity (mPa.s)	1500	570	280	20
-------------------	------	-----	-----	----

**Example 6: Interactions between guar gum and xanthan gum.**

The compositions were prepared as in Example 1. The viscosity was measured with a Brookfield viscosimeter (spindle LV) at 20°C.

**A. Effects on the viscosity.**

Xanthan gum (g)	0	0.3	0.9	1.5	2.1
Guar gum (g)	3	2.7	2.1	1.5	0.9
Water (g)	197	197	197	197	197
Viscosity (mPa.s)	49000	212000	60000	6000	2100

**B. Influence of collagen 5% by weight on the viscosity.**

Xanthan gum (g)	0	0.3	0.9	1.5	2.1
Guar gum (g)	3	2.7	2.1	1.5	0.9
Collagen (g)	10	10	10	10	10
Water (g)	187	187	187	187	187
Viscosity (mPa.s)	100	15000	98000	200000	300000

**C. Influence of acidity and collagen 5% by weight on the viscosity.**

Xanthan gum (g)	0	0.3	0.9	1.5	2.1
Guar gum (g)	3	2.7	2.1	1.5	0.9
Collagen (g)	10	10	10	10	10
Citric acid (g)	0.05	0.05	0.05	0.05	0.05
Water (g)	187	187	187	187	187
Viscosity (mPa.s)	100	1200	200000	320000	340000

**D. Influence of acidity and collagen 2.5% by weight on the viscosity.**

Xanthan gum (g)	0	0	0.3	0.3	0.9	0.9	0.9
Guar gum (g)	3	3	2.7	2.7	2.1	2.1	2.1
Collagen (g)	5	5	5	5	5	5	5
Citric acid (g)	0.025	0.25	0.025	0.25	0.025	0.25	1
Water (g)	198	198	198	198	198	198	197
Viscosity (mPa.s)	100	5000	1200	25000	200000	42600	39000

**Example 7: Preparation of a lemon instant drink containing guar gum and collagen.**

The different ingredients are admixed together and mixed with 264.4 g of water at temperature of about 25°C, in a shaker, for about 10 seconds.

Guar gum <sup>(1)</sup> (g)	3.75
Collagen (g)	17.3
Fructose (g)	12.3
Na <sub>2</sub> Hpyrophosphate (g)	0.9
Citric acid (g)	0.5
Lemon flavor (g)	0.7
Aspartame <sup>(2)</sup> (g)	0.015
Acesulfam K <sup>(3)</sup> (g)	0.015
Acetic ester of mono and diglycerides (g)	0.08

<sup>(1)</sup> The guar gum is of high molecular weight type.

The final composition has a pH of 4.35 and a viscosity from 25 to 30 mPa.s.

**Claims**

1. A composition comprising one or more viscous soluble fiber(s) and one or more viscosity-lowering protein(s).
2. A composition according to claim 1 wherein said protein has a molecular weight of less than 30 kDa.
3. A composition according to claim 1 or claim 2 wherein said protein is selected from the group consisting of wheat protein, egg protein, collagen, whey protein, casein, soy protein, pea protein, muscle protein, gluten, fibrillar protein, silk protein, hydrolysates thereof and mixtures thereof.
4. A composition according to any preceding claim wherein said fiber is selected from the group consisting of gum, mucilage, carrageenan, pectin, beta-glucan, and mixtures thereof.
5. A composition according to claim 4 wherein said gum is selected from the group consisting of konjac gum, locust bean gum, tara bean gum, gum tragacanth, karaya gum, xanthan gum, arabic gum, guar gum, gellan gum and mixtures thereof.
6. A composition according to claim 5 wherein said gum comprises guar gum.
7. A composition according to any preceding claim wherein said fiber comprises guar gum and one or more viscous soluble fiber(s) selected from the group consisting of pectin, beta-glucan, xanthan gum, arabic gum and konjac gum.
8. A composition according to any preceding claim wherein said composition has a viscosity lower than 500 mPa.s at room temperature.
9. A composition according to any preceding claim wherein said composition comprises about 1% to about 5% by weight of viscous soluble fibers, based on the total weight of the composition.
10. A composition according to any preceding claim wherein the ratio of said fiber to said protein is

11. A composition according to any preceding claim wherein said composition is non-viscous.
12. A composition according to any preceding claim wherein said composition is ready-to-consume.
13. A composition according to claim 12 wherein said composition comprises about 1% to about 60% by weight of viscous soluble fibers, based on the total weight of the composition.
14. A composition according to any one of claims 1 to 13 for use as a medicament.
15. Use of a composition according to any one of claims 1 to 13 in the manufacture of a medicament or nutritional composition for the prevention or treatment of metabolic syndrome and/or diabetes and/or for lowering blood cholesterol level.
16. A method of treating and/or preventing metabolic syndrome and/or diabetes, and/or of lowering blood cholesterol level in a mammal in need of such a treatment comprising administering to said mammal an effective amount of a composition according to any one of claims 1 to 13.
17. A method for providing viscous soluble fibers to a mammal in need of such fibers comprising administering to said mammal a composition according to any one of claims 1 to 13.
18. Use of a protein to modulate the viscosity of a composition comprising one or more viscous soluble fiber(s), wherein said protein is selected from the group consisting of wheat protein, egg protein, collagen, whey protein, casein, soy protein, pea protein, muscle protein, gluten, fibrillar protein, silk proteins, hydrolysates thereof and mixtures thereof.
19. Use of a protein according to claim 18, wherein said viscous soluble fiber is selected from the group consisting of gum, pectin, beta-glucan, mucilage, carrageenan, xanthan gum, arabic gum, konjac gum and mixtures thereof.
20. Use of a protein according to claim 19, wherein said gum is selected from the group consisting of konjac gum, locust bean gum, tara bean gum, gum tragacanth, karaya gum, xanthan gum, arabic gum, guar gum, gellan gum

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**